

AALENIAN *TMETOCERAS* (AMMONOIDEA) FROM IBERIA

Taphonomy and Palaeobiogeography

S. R. Fernández-López,¹ M. H. Henriques,² A. Linares,³ J. Sandoval,³ and
M. S. Ureta¹

¹Dept. y UEI de Paleontología
Facultad de Ciencias Geológicas (UCM) e Instituto de Geología Económica
(CSIC-UCM)

28040-Madrid, Spain

²Dept. Ciências da Terra

Centro de Geociências

Universidade de Coimbra

3049-Coimbra Codex, Portugal

³Dept. de Estratigrafía y Paleontología

Facultad de Ciencias

Universidad de Granada

18071 Granada, Spain

Abstract

From different areas of the Iberian Peninsula more than 600 specimens of Aalenian *Tmetoceras* have been found. This taxonomic group represents less than 20% of the whole ammonoids recorded in Opalinum, Murchisonae, Bradfordensis and Concavum biozones. *Tmetoceras* representatives, as well as Phylloceratina and Lytoceratina, were more frequent in shelfal basins than in epicontinental platforms. Taphonomic data suggest a eudemic character of the representatives of *T. scissum* in shelfal basins or oceanic areas. Exceptional immigrants and drifted shells of this species arrived in shallow environments of neighbouring platforms. In contrast, representatives of *T. regleyi* inhabited preferentially shallow environments of epicontinental platforms.

T. scissum was a pandemic species, inhabiting oceanic or shelfal environments in the early Aalenian. However, some species of *Tmetoceras*, such as *T. regleyi* and *T. flexicostatum*, were geographically restricted in very distant areas. *T. regleyi* has been found only in Euro-

pean areas of the West Tethyan Subrealm. A pattern of adaptive radiation may have taken place in the Western Tethys during the Opalinum-Murchisonae biochrons, giving rise to *T. regleyi* from *T. scissum*. Specialized forms of *Tmetoceras* (k-strategists such as the individuals of the species *T. regleyi*) are widespread in the epicontinental platforms around the Western Tethys during the Murchisonae and Bradfordensis biochrons. Epicontinental, specialized forms of *T. regleyi* suffered extinction in the latest Bradfordensis Biochron. Shelfal or oceanic, generalist forms of *T. scissum* disappeared in the Western Tethys or the Mediterranean Province in the latest Bradfordensis Biochron, but they survived in the East-Pacific Subrealm.

1. INTRODUCTION

The subfamily Tmetoceratinae is a monogeneric group, *Tmetoceras* (M+m), comprising several European forms distinguished at specific level on the basis of morphological differences: *T. scissum* (Benecke, 1865), *T. regleyi* (Dumortier, 1874), *T. difalense* (Gemmellaro, 1886) and *T. hollandae* (Buckman, 1892). However, when fossils are abundant, these species all appear to intergrade (cf. Callomon and Chandler, 1994).

The main purpose of this paper is to show that the distribution of *Tmetoceras* shells around the margins of the Iberia during the Aalenian was influenced by biogeographical and taphonomic dispersal, related to changes in the relative depth of sea level.

2. TAPHONOMIC CHARACTERS AND DISTRIBUTION OF AALENIAN TMETOCERAS IN IBERIA

More than 600 specimens of Aalenian *Tmetoceras* found *in situ* in different areas of the Iberian Peninsula have been studied (Figure 1). These specimens have been found in the following areas and outcrops:

1. N Lusitanian Basin (Cabo Mondego, Coimbra and Degracias; cf. Perrot, 1955; Mouterde *et al.*, 1964, 1980; Rocha *et al.*, 1981; Henriques, 1989, 1992, 1995).
2. Asturias (Santa Mera and El Puntal; cf. Suárez-Vega, 1974; Fernández-López and Suárez-Vega, 1980).
3. Basque-Cantabrian Basin (Camino, Castillo Pedroso, Cillamayor, Rebolledo de la Torre and San Andrés; cf. Fernández-López *et al.*, 1988b; Goy *et al.*, 1990; Canales *et al.*, 1993).
4. NE Cameros (Préjano and Muro de Aguas; cf. Goy and Ureta, 1981; Ureta, 1985, 1988).
5. NW Iberian Basin (Canales de la Sierra, Castrovido, Hontoria, Rabanera, Talveila and Villavelayo; cf. Ureta, 1985; Ureta and Goy, 1986; Fernández-López *et al.*, 1988a).
6. Aragonese Platform (Obón and Ricla; cf. Mouterde *et al.*, 1978; Fernández-López and Aurell, 1988; Goy *et al.*, 1988).
7. Tortosa Platform (Coll de Soms, Mas de Rojals, Vall Llarga and Vandellós; cf. Fernández-López and Mouterde, 1985; Fernández-López *et al.*, 1996).
8. NW Castilian Platform (Arcos de Jalón and Fuentelsaz; cf. Goy and Ureta, 1987, 1991; Goy *et al.*, 1994).
9. Central Castilian Platform (Barranco de la Hontanilla, Peracense and Rambla del Salto; cf. Fernández-López and Gómez, 1990a, b).


Figure 1. Areas of Iberia mentioned in text: 1) N Lusitanian Basin, 2) Asturias, 3) Basque-Cantabrian Basin, 4) NE Cameros, 5) NW Iberian Basin, 6) Aragonese Platform, 7) Tortosa Platform, 8) NW Castilian Platform, 9) Central Castilian Platform, 10) SE Castilian Platform, 11) Majorca, 12) External Subbetic Basin, 13) Oriental sector of Median Subbetic Basin, 14) Central sector with condensed sections of Median Subbetic Basin, and 15) Central sector with expanded sections of Median Subbetic Basin.



10. SE Castilian Platform (Ribarroja and Domeño; cf. Fernández-López, 1980, 1985).
11. Majorca (Cúber and Llodrá; cf. Goy and Ureta, 1988; Álvaro *et al.*, 1989).
12. External Subbetic Basin (Cortijo Veteta and Sierra de Gaena; cf. Sandoval, 1983).
13. Oriental sector of Median Subbetic Basin (Sierra de Ricote; cf. Seyfried, 1978; Linares and Sandoval, 1993; García-Gómez *et al.*, 1994).
14. Central sector with condensed sections of Median Subbetic Basin (Cerro Méndez and Río Fardes; cf. Linares, 1979; Hernández-Molina *et al.*, 1991; Linares and Sandoval, 1993; García-Gómez *et al.*, 1994).
15. Central sector with expanded sections of Median Subbetic Basin (Cerro Méndez, Montillana and Sierra de San Pedro; cf. García-Gómez *et al.*, 1994; Linares, 1979; Linares *et al.*, 1988; Linares and Sandoval, 1992, 1993).

These areas of the Iberian Peninsula comprise two separate habitats: epicontinental platforms and shelfal basins. Habitats of epicontinental platform were cratonic areas flooded by shallow marine waters. In contrast, habitats of shelfal basins were situated on the continental shelf showing open marine and oceanic environments.

Two shelfal basins were located in Tethyan and Protoatlantic margins of the Iberian Plate throughout Aalenian time: the Betic and the Lusitanian basins (Figure 2). A set of very shallow epicontinental platforms, the so-called Iberian Basin, was situated between them. Depositional environments of these Aalenian basins varied from those of very shallow outer platform to those of deep basin. Aalenian strata are fundamentally limestones which may attain 85 m thick (Henriques *et al.*, 1996). Ammonoid assemblages of these environments varied from those with a predominance of *Phylloceratina* and *Lytoceras* to those that were mainly *Ammonitina*. For example, ammonoid assemblages in the Median



Subbetic Basin are dominated by Phylloceratina and Lytoceratina during the Bradfordensis Biochron. In Lusitanian and Iberian basins, Ammonitina are the most frequent Aalenian ammonoids found, and Phylloceratina or Lytoceratina are very scarce. Fossils of *Tmetoceras* are also more abundant in the Betic Basin than in the Lusitanian or in the Iberian basins (Figure 3). The subfamily Tmetoceratinae represents less than 20% of the whole ammonoids recorded in Opalinum, Murchisonae, Bradfordensis, and Concavum biozones. However, from a taphonomic and palaeoecological point of view, the distribution of Aalenian *Tmetoceras* shows conspicuous variations around the Iberian Plate at a zonal scale (Tables 1-4).

Fragmentary, incomplete specimens are commonly represented among Aalenian ammonoids of the Iberian Peninsula. No accumulated *Tmetoceras* have been identified among them. Fossil assemblages show high degrees of removal (*i.e.*, proportion of resedimented or reelaborate elements). The palaeontological material from Asturias, Basque-Cantabrian and Iberian basins is composed of resedimented or reelaborate elements. They represent individuals with less than 50 mm in diameter. However, Betic and Lusitanian *Tmetoceras* are mostly composed of resedimented specimens showing diameter values between 13 and 78 mm.

Two species of *Tmetoceras* have been recognized: *T. scissum* and *T. regleyi* (cf. Fernández-López *et al.*, 1997). On the basis of morphological criteria, *T. scissum* comprise evolute shells having thin, acute, prominent and distant ribbing, and they have a moderately complex suture line (Fig. 3A-D, F-G). In contrast, representatives of *T. regleyi* are more involute, showing thick, blunt and dense ribs, and their suture line is more simple (Figure 3E).

Specimens of *T. scissum* are more abundant and they have higher stratigraphical persistence in the Betic and Lusitanian basins than in the epicontinental platforms of Iberia (Figure 4). In contrast, specimens of *T. regleyi* are practically restricted to areas of epicontinental platform, where they are the dominant species of the genus during the Murchisonae and Bradfordensis biochrones. Nevertheless, very scarce adult specimens of *T. regleyi* have been found in the Murchisonae and Bradfordensis biozones from the Betic Basin.

Shells of *Tmetoceras* recorded in Aalenian deposits of these areas compose taphonic populations of three types: TP1, TP2 and TP3 (Figure 5). Taphonic populations of type 1

Figure 3. Aalenian *Tmetoceras* of the Iberian Peninsula. (A) Juvenile individuals of *Tmetoceras scissum*. Resedimented elements with complete body chambers. CMB.19.1. Opalinum Biozone. Cerro Méndez (Central sector with expanded sections of Median Subbetic Basin). x2. (B) Post-juvenile individuals of *Tmetoceras scissum*. Resedimented elements with incomplete body chambers. JRI.1.63.2. Opalinum Biozone. Sierra de Ricote (Oriental sector of Median Subbetic Basin), x1. (C) Adult individual of *Tmetoceras scissum*. Reelaborated element with complete body showing a disarticulation surface (DS). CM.7.75. Murchisonae Biozone. Cerro Méndez (Central sector with condensed sections of Median Subbetic Basin), x1. (D) Immature individual of *Tmetoceras scissum*. Reelaborated element with incomplete body chamber. CM.A5.20. Murchisonae Biozone. Cerro Méndez (Central sector with condensed sections of Median Subbetic Basin), x1. (E) Post-juvenile individual of *Tmetoceras regleyi*. Reelaborated element with incomplete body chamber. 5SS126/14. Bradfordensis Biozone. Coll de Soms (Tortosa Platform), x1. (F) Post-juvenile individual of *Tmetoceras scissum*. Reelaborated element with incomplete body chamber. PV12/23. Bradfordensis Biozone. Ribarroja (SE Castilian Platform), x1. (G) Post-juvenile individual of *Tmetoceras scissum*. Resedimented element with incomplete phragmocone. SG.107.1. Opalinum Biozone. S. Guião, Degraças (N Lusitanian Basin), x1. (H) Internal mould of a shell of ammonites (*Brasilio*) showing traces of open and lobulate fractures (F) likely produced by crustaceans. Bradfordensis Biozone. Domeño (SE Castilian Platform), x1.




Table 1. Characters of the Opalinum *Tmetoceras* in different areas of the IberiaOpalinum
Biozone

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
	NE LYCERATINA BASIN	ASTONIA	BAJONA-LANZAROTE BASIN	SE CANTABRIA	SE BAYONA BASIN	ANDORRITA PLATON	TORTOSA PLATON	SE CANTABRIA PLATON	CENTRAL CANTABRIA PLAT	SE CANTABRIA PLATON	MADRUGA	ESTREMA TROMBIC BASIN	MEDIO TROMBIC BASIN (TROMBIC MOUNTAIN)	MEDIO TROMBIC BASIN (TROMBIC MOUNTAIN)	MEDIO TROMBIC BASIN (TROMBIC MOUNTAIN)
Specimens of <i>Tmetoceras</i>	>34	>18	23	14	22	4	3	3	1	24	3	6	37		150
Frequency	17	20	16	18	12	<1	<1	<1	<1	24		13	15		21
Fossiliferous levels	34	7	15	5	14	4	3	3	1	1	3	1	5		46
Degree of packing		61	35	64	36	0	0	0	0	90	0	83	86		69
Degree of taphonomic heritage	0	0	0	0	22	0	0	0	0	80	0	85	0		0
Incomplete phragmocones (%)	60	98	69	36	37	100	100	100	100	37	33	0	0		0
Size-frequency distribution															
uni- (U) or polymodal (P)	U	U	U		U					U			U		U
asymmetric (A) or normal (N)	N	A	A		A					A			N		A
negative (-) or positive (+) skew			+		+					+			+		+
Shells of juvenile individuals															
absent (A) rare (R) predominant (P)	P	R		N	N					P	P	R			R
Shells of adult individuals															
absent (A) rare (R) predominant (P)	N	N	R			P	P	P		R		P	R		
Proportion of elements/species	>34	>18	23	14	11	4	3	3	1	20	3	6	37		
Diversity of the genus	1	1	1	1	2	1	1	1	1	1	1	1	1		1
Type of taphonic population	TP1-2	TP1-2	TP1-2	TP1-2	TP1-2	TP3	TP3	TP3	TP3	TP1	TP1-2	TP2	TP1-2		TP1-2
Frequency of <i>Phylloceratina</i>	0	0	0	0	0	<1	0	0	0	0		>10	14		13
Frequency of <i>Lytoceratina</i>	<1	<1	0	0	0	0	0	0	0	0		>5	4		5

Table 2. Characters of the Murchisonae *Tmetoceras* in different areas of the IberiaMurchisonae
Biozone

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
	NE LYCERATINA BASIN	ASTONIA	BAJONA-LANZAROTE BASIN	SE CANTABRIA	SE BAYONA BASIN	ANDORRITA PLATON	TORTOSA PLATON	SE CANTABRIA PLATON	CENTRAL CANTABRIA PLAT	SE CANTABRIA PLATON	MADRUGA	ESTREMA TROMBIC BASIN	MEDIO TROMBIC BASIN (TROMBIC MOUNTAIN)	MEDIO TROMBIC BASIN (TROMBIC MOUNTAIN)	MEDIO TROMBIC BASIN (TROMBIC MOUNTAIN)
Specimens of <i>Tmetoceras</i>			5					1				1	13	78	
Frequency			9					<1					2	18	14
Fossiliferous levels			3					1					1	3	23
Degree of packing			40					0					0	77	70
Degree of taphonomic heritage			0					0					0	100	0
Incomplete phragmocones (%)			60					100					0	0	0
Size-frequency distribution															
uni- (U) or polymodal (P)															U
asymmetric (A) or normal (N)															A
negative (-) or positive (+) skew															+
Shells of juvenile individuals															
absent (A) rare (R) predominant (P)			N												P
Shells of adult individuals															
absent (A) rare (R) predominant (P)															
Proportion of elements/species			5					1					1	13	39
Diversity of the genus			1					1					1	1	2
Type of taphonic population			TP2					TP3					TP3	TP2	TP1
Frequency of <i>Phylloceratina</i>			0					0					12	30	12
Frequency of <i>Lytoceratina</i>			5					0					>6	9	3

Table 3. Characters of the Bradfordensis *Tmetoceras* in different areas of the Iberia

**Bradfordensis
Biozone**

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
	NE LUSITANIAN BASIN	ASTURIA	NEOGENE CANTABRIAN BASIN	NE CANTABRIA	NE IBERIAN BASIN	NEOGENE PLATONOME	TORTOSA PLATONOME	NE CANTABRIAN PLATONOME	CANTABRIAN PLATONOME	NE CANTABRIAN PLATONOME	MAJADITA	EXTREMADURA TERTIARY BASIN	MURCIA TERTIARY BASIN	MURCIA TERTIARY BASIN	MURCIA TERTIARY BASIN
Specimens of <i>Tmetoceras</i>	»30	+		2	+		34	3	2	25			1	1	75
Frequency	17	+		2	+		3	«1	«1	17			2	10	10
Fossiliferous levels	15			1			3	2	2	1			1		29
Degree of packing	»50	+		50	+		97	33	0	96			0	0	61
Degree of taphonomic heritage	20			0	+		0	0	50	100			0	100	0
Incomplete phragmocones (%)	80	+		100			42	100	100	20			0	0	0
Size-frequency distribution															
uni- (U) or polymodal (P)							U	-	-	U					U
asymmetric (A) or normal (N)							A			A					A
negative (-) or positive (+) skew							-			+					-
Shells of juvenile individuals															
absent (A) rare (R) predominant (P)		+		+	+		P	+	-	P			+		P
Shells of adult individuals															
absent (A) rare (R) predominant (P)	R	+		-	-		R	P	P	R			+		-
Proportion of elements/species	»30			2	+		34	1	2	25			1	1	37
Diversity of the genus	1			1			1	1	1	1			1	1	2
Type of taphonic population	TP1-2			TP1-2			TP1	TP3	TP3	TP1			TP3	TP3	TP1
Frequency of <i>Phylloceratina</i>	0	0	0	0	0	«1	0	0	0	0			8	60	17
Frequency of <i>Lytocerotina</i>	3	«1	0	0	0	0	0	0	0	0			4	20	8

Table 4. Characters of the Concavum *Tmetoceras* in different areas of the Iberia

**Concavum
Biozone**

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
	NE LUSITANIAN BASIN	ASTURIA	NEOGENE CANTABRIAN BASIN	NE CANTABRIA	NE IBERIAN BASIN	NEOGENE PLATONOME	TORTOSA PLATONOME	NE CANTABRIAN PLATONOME	CANTABRIAN PLATONOME	NE CANTABRIAN PLATONOME	MAJADITA	EXTREMADURA TERTIARY BASIN	MURCIA TERTIARY BASIN	MURCIA TERTIARY BASIN	MURCIA TERTIARY BASIN
Specimens of <i>Tmetoceras</i>						2		2	+						
Frequency						«1		«1							
Fossiliferous levels						2		2	+						
Degree of packing						0		0	+						
Degree of taphonomic heritage						100		100	+	+					
Incomplete phragmocones (%)						100		50							
Size-frequency distribution															
uni- (U) or polymodal (P)						+									
asymmetric (A) or normal (N)						+									
negative (-) or positive (+) skew						-									
Shells of juvenile individuals															
absent (A) rare (R) predominant (P)															
Shells of adult individuals															
absent (A) rare (R) predominant (P)						P		P	+	+					
Proportion of elements/species						2		2	+	+					
Diversity of the genus						1		1							
Type of taphonic population						TP3		TP3							
Frequency of <i>Phylloceratina</i>	0	0	0	0	0	0	0	0	0	0	0		10	30	»5
Frequency of <i>Lytocerotina</i>	2	0	0	0	0	0	0	0	0	0	0		0	3	2

	Protoatlantic (1)	Iberia (2)(3)(4)(5)(6)(7)(8)(9)(10)	W Tethys (12)(13)(14)(15)
	Shelfal basin	El continental platform	Shelfal basin
Concavum Bc.	0	4	0
	0 0	4 0	0 0
Bradfordensis Bc.	30	63	77
	30 0	27 36	72 5
Murchisonae Bc.	0	6	92
	0 0	0 6	88 4
Opalinum Bc.	34	112	193
	34 0	110 2	193 0
Total number of <i>Tmetoceras</i> = 611			
<i>T. scissum</i> = 558 <i>T. regleyi</i> = 53			

Figure 4. Total number of Aalenian *Tmetoceras* obtained from each basin, indicating the specific number of specimens of *T. scissum* and *T. regleyi*.

(TP1) are composed of monospecific shells showing unimodal and asymmetric distribution of size-frequencies, with positive skew. These populations have a high proportion of microconchs and the shells of juvenile individuals are predominant, whilst shells of adult individuals are scarce. Taphonic populations of type 2 (TP2) are composed of mono- or polyspecific shells showing unimodal and normal distribution of size-frequencies, with high kurtosis. Populations of this second type have a low proportion of microconchs and the shells of juvenile individuals are scarce, whilst the shells of adult individuals are common. Taphonic populations of type 3 (TP3) are composed of polyspecific shells showing uni- or polymodal and asymmetric distribution of size-frequencies, with negative skew. Shells of juvenile individuals are absent, microconchs are very scarce and shells of adult individuals are predominant in taphonic populations of this last type.




Figure 5. Types of taphonic populations of ammonites developed in environments of shallow outer platform as observed on Middle Jurassic ammonites of the Iberian Basin (after Fernández-López, 1995).




Figure 6. Chart showing the different values of packing degree (PD) and taphonomic heritage degree (THD) in the different areas, at a zonal scale, indicating their type of taphonic population (TP-1, TP-2, TP-3).

Different types of taphonic populations of *Tmetoceras* show also distinctive degrees of packing (Figure 6). This index is directly proportional to the abundance of specimens and it takes higher values when the number of fossiliferous levels decreases. Taphonic populations of type 1 have high values of packing (up to 55%). Taphonic populations of type 3 have low values of packing (under 30%). Thus, the degree of packing is another criterion to determine the type of taphonic population. In contrast, the degree of taphonomic heritage, *i.e.* the proportion of reelaborated elements, is not correlated with the type of taphonic population but with the degree of stratigraphical or sedimentary condensation. Taphonic populations of type 1 showing very high degrees of packing and taphonomic heritage were formed in condensed sections from the Median Subbetic Basin during the Murchisonae Biochron, as well as in condensed associations from the External Subbetic Basin during the Opalinum Biochron and from SE Castilian Platform during the Opalinum and Bradfordensis biochrons. Taphonic populations of type 3 showing very high degrees of taphonomic heritage associated with very low degrees of packing were formed in condensed sections from the Median Subbetic Basin during the Bradfordensis Biochron. Some taphonic populations of type 3 displaying very high degrees of taphonomic heritage occur also in condensed associations from Aragonese Platform during the Concaum Biochron and from NW Castilian Platform during the Bradfordensis Biochron.

Opalinum *Tmetoceras* are fairly common in Lusitanian, Iberian and Betic basins (Figure 7). Their frequencies are lower than 1% in the Tortosa, Aragonese and Castilian Platforms. Taphonic populations developed in the Betic and Lusitanian basins, as well as in the deepest environments of the Iberian Basin, were of types 1 and 2. In contrast, taphonic populations were of type 3 in the shallower environments of the Iberian Basin. All the Opalinum *Tmetoceras* found in the Betic and Lusitanian basins are *T. scissum* (Figure 8). However, a post-juvenile specimen of *T. regleyi* has been found in the upper portion of the Opalinum Biozone, associated with abundant specimens of *T. scissum* in the NW Iberian Basin.




Figure 7. Frequency of *Tmetoceratinae* and *Phylloceratina* plus *Lytoceratina* in different areas of Iberia, and some size-frequency distributions of *Tmetoceras*, from the Opalinum Biozone.

In the Iberian Peninsula, Murchisonae rocks are thin compared with the Bradfordsen rocks above or with the Opalinum below. Murchisonae ammonites are also more scarce. Specimens of *Tmetoceras scissum* are frequent in some Betic outcrops only, and they correspond mainly to monospecific taphonic populations of type 1 (Figure 9). *Tmetoceras* recorded in the Murchisonae Biozone of the Basque-Cantabrian Basin comprise local polyspecific assemblages (Figure 10). A few specimens of *Tmetoceras scissum* have been obtained from the Basque-Cantabrian Basin, associated with scarce juvenile and post-juvenile specimens of *T. regleyi* that represent probable demic individuals. Some specimens of *T. scissum* founded in the NW areas of the Castilian Platform are adult. A few adult specimens of *Tmetoceras regleyi* have been obtained also from the central sector of Median Subbetic Basin, associated with abundant specimens of *T. scissum*. No Murchisonae *Tmetoceras* are known from the Lusitanian Basin.




Figure 8. Frequency of Opalinum *Tmetoceras*, indicating the type of taphonic population of *T. scissum* or *T. regleyi* in each area.




Figure 9. Frequency of *Tmetoceratinae* and *Phylloceratina* plus *Lytoceratina* in different areas of Iberia, and the size-frequency distribution of *Tmetoceras* specimens in the central sector with expanded sections of Median Subbetic Basin, from the Murchisonae Biozone.

Ammonoids are more frequent and widespread in the Bradfordensis Biozone than in the Murchisonae Biozone of Iberia. Bradfordensis *Tmetoceras* are quite frequent in Median Subbetic Basin, Lusitanian Basin and SE Castilian Platform (Figure 11). Taphonic populations of *T. scissum* developed in the Betic and Lusitanian basins, as well as in the SE Castilian Platform, were of types 1 and 2 (Figure 12). In contrast, taphonic populations of *T. scissum* were of type 3 in shallower areas of the Iberian Basin and Betic Basin. However, specimens of *T. regleyi* become relatively frequent in some deep areas of the Iberian Basin, composing taphonic populations of type 1, whilst in the Betic Basin only very scarce adult specimens of this species have been found.

Bradfordensis taphonic populations of *T. scissum* in the SE Castilian Platform, as well as taphonic populations of *T. regleyi* in the Tortosa Platform, show distinctive features (Figure 11). These populations are of type 1 and they display higher proportions of incomplete phragmocones among the largest specimens. These higher proportions of fragmentary shells among the largest specimens is indicative of fragmentation of the shells by predators or scavengers associated with low values of sedimentation rates. Under opposite conditions, with high values of sedimentation rates and showing no evidence of predators or scavengers, the mechanical fragmentation of shells by turbulence produced higher proportions of incomplete phragmocones among the smallest specimens (Fernández-López and Mouterde, 1994; Fernández-López, 1997). The common occurrence of open and lobulate fractures in ammonite shells of these two areas also confirm this interpretation (Figure 3H).




Figure 10. Frequency of Murchisonae *Tmetoceras*, indicating the type of taphonic population of *T. scissum* or *T. regleyi* in each area.




Figure 11. Frequency of *Tmetoceratinae* and *Phylloceratina* plus *Lytoceratina* in different areas of Iberia, and some size-frequency distributions of *Tmetoceras*, from the Bradfordensis Biozone.

Several *Tmetoceras* have been obtained from condensed associations along the Iberian Cordillera in deposits of the upper Aalenian and of the lower Bajocian (Figures 13, 14). These specimens are the only *Tmetoceras* identified in the Iberian Plate in post-Bradfordensis sediments, but they are reelaborate fossils. No Concavum *Tmetoceras* are known from the Lusitanian or Betic basins.

3. PALAEOECOLOGICAL INTERPRETATIONS AND HABITATS

Type of taphonic population, abundance and frequency of specimens, stratigraphic persistence and degree of packing are distinctive criteria in interpreting the palaeoecology and palaeobiogeography of the recorded associations of ammonites. Biogeographical dispersal and necroplanktic drift of shells (a case of taphonomic dispersal) were the main determining factors on the final distribution of the ammonite shells (Figure 15). Consequently, the present distribution of ammonite shells reflect both their original and post-mortem distribution.




Figure 12. Frequency of *Bradfordensis Tmetoceras*, indicating the type of taphonic population of *T. scissum* or *T. regleyi* in each area.




Figure 13. Frequency of *Tmetoceratinae* and *Phylloceratina* plus *Lytoceratina* of the Concavum Biozone in different areas of Iberia.

In Lusitanian and Iberian basins, representatives of *Ammonitina* are dominant among the Aalenian ammonoids, and the scarce specimens of *Phylloceratina* or *Lytoceratina* represent ademic organisms. In Betic Basin, as an area of the West Tethys, Aalenian *Phylloceratina* and *Lytoceratina* were represented by endemic organisms.

Eudemic *Tmetoceras* are represented in the fossil record by taphonic populations of type 1, characterized by the relative abundance of juvenile elements and high values of taphonomic packing. In contrast, ademic *Tmetoceras* are represented by taphonic populations of type 3, showing low values of taphonomic packing, and they are dominated by adult elements. The active and passive biotransport led to the occurrence of immigrant and transported *Tmetoceras*, which respectively are midemic and parademic organisms. Bi-dispersed *Tmetoceras* are represented in the fossil record by taphonic populations of type 2 or 3 showing median values of taphonomic packing.

Taphonomic data suggest a eudemic character to the *T. scissum* representatives recorded in shelfal basins of the Iberian Plate (Figure 16). Aalenian *T. scissum* are widely distributed in the Betic and Lusitanian basins, where they occur in rich but nearly monospecific associations. Eudemic *T. scissum* inhabited basin areas, characterized by the occurrence of juvenile specimens and taphonic populations of type 1. In contrast, many allochthonous elements of *T. scissum* from the Iberian Basin are interpreted as having arrived at their present locations by neoproplanktic drift. The taphonomic dispersal produced the concentration of adult shells of *T. scissum* in shallow neritic environments, representing ademic organisms. They did not breed or ontogenetically develop in the Castilian and Aragonese platforms. In the Castilian Platform, there is evidence of shell transport from the north (Fernández-López and Gómez, 1990a). Northwestwards from Castilian and Aragonese platforms, the NW Iberian Basin, NE Cameros, Basque-Cantabrian Basin and Asturias were the depositional sites of autochthonous and allochthonous shells of *T. scissum* throughout the Aalenian time.




Figure 14. Frequency of Concavum *Tmetoceras*, indicating the type of taphonic population of *T. scissum* in each area.




Figure 15. Summary of the palaeobiogeographical categories mentioned in this work. The palaeobiological entities, in particular ammonites, are demic when their remains are found in their living area. Conversely, they are ademic organisms or taxa when they are inferred outside of their living area. Demic species may be inferred in their breeding area (eudemic), in an area normally occupied but where they do not breed (miodemic), or in a living area occasionally reached by passive bio-dispersal (parademic; after Fernández-López, 1990, 1991 and Fernández-López and Meléndez, 1995).

Some Opalinum specimens exhibiting morphological characters of *T. regleyi* have been found in the NW Iberian Basin, but they are interpreted as post-juvenile, ademic organisms (Figure 17). The active and passive biodispersal led to the occurrence of immigrant and transported *T. regleyi* in the Iberia, which respectively represent miodemic and parademic organisms. The active biodispersal led to the occurrence in outer platforms of immigrant *T. regleyi*, which represent miodemic organisms, during the Murchisonae Biochron. Taphonic populations of *T. regleyi* dominated by juvenile individuals were produced by eudemic organisms in the Iberian Basin during the Bradfordensis Biochron. Scarce adult specimens of *T. regleyi* confined to isolated horizons in the Betic Basin are interpreted as evidence of parademic organisms dispersed during the Murchisonae and Bradfordensis biochrons.

In conclusion, shells of Aalenian *Tmetoceras* have been accumulated in widely distributed areas throughout the Iberian Plate during the Opalinum Biochron. However, the taphonomic and biogeographical dispersal of *Tmetoceras* through the Iberian Basin was hindered during the Murchisonae Biochron, as a consequence of the relative fall of sea level. A new relative rise of sea level during the Bradfordensis Biochron favoured the immigration of juvenile *Tmetoceras* in the deepest areas of the epicontinental platforms. Se-




Figure 16. Zonal chart of the palaeobiogeographical distribution of *Tmetoceras scissum* representatives in the Iberia.




Figure 17. Zonal chart of the palaeobiogeographical distribution of *Tmetoceras regleyi* representatives in the Iberia.

veral specimens of *Tmetoceras* have been found in the rocks of the Concavum Biozone, but they are reelementary elements derived from older sediments. The Iberian Basin was not active as an epicontinental migration route for Aalenian *Tmetoceras*. *T. scissum* inhabited environments of shelfal basins. *T. regleyi* inhabited environments of epicontinental platforms. However, *T. scissum* occurs together with *T. regleyi* in beds of the Iberian and Betic basins. These two species appear to intergrade, but they were ecological and biogeographically differentiated. Several lines of taphonomic evidences (such as type of taphonic population, abundance, frequency, stratigraphic persistence, degree of packing, and degree of taphonomic heritage) support the concept that the environments of outer epicontinental platform were the sites inhabited by *T. regleyi*.

4. GLOBAL DISTRIBUTION OF TMETOCERAS

Tmetoceras can be considered a pandemic genus of ammonites abundantly represented in the northern margins of the West Tethyan Subrealm, from north Africa and western Europe eastwards to southeast Asia and beyond, and around the Pacific Ocean, including the East-Pacific Subrealm (Figure 18). Although no abundant Aalenian *Tmetoceras* are known from the Arctic Province, some species have been found in Alaska (Westermann, 1964, 1981, 1992). Aalenian *Tmetoceras* occur in the following two provinces of the Boreal Realm: the Bering Province (North Pacific margin and northwestern North America; cf. Imlay, 1955, 1962, 1964; Poulton, 1982; Poulton and Tipper, 1991; Poulton *et al.*, 1991) and the Sub-Boreal Province (Scotland, northern France, northern Germany, Poland; Morton, 1965; Bielecka *et al.*, 1970; Morton and Hudson, 1995).

Specimens of *T. scissum* have been found in the following three provinces of the West Tethyan Subrealm:

1. Northwest European Province (northwestern areas of the Iberian Basin, Britain, northern France, Germany, Rumanian Carpathians; cf. Buckman, 1892, 1907; Arkell, 1933; Patruşius and Popa, 1971; Dietl, 1977; Dietl and Etzold, 1977; Callomon and Chandler, 1990, 1994; Callomon, 1995).



Figure 18. Palaeogeographical distribution of *Tmetoceras* representatives.

2. Sub-Mediterranean Province (epicontinental seas of North Africa, northern Lusitanian Basin, southeastern areas of the Iberian Basin, southern and central France, southern Germany, Switzerland, Iran, Caucasus; cf. Flamand, 1911; Fischer, 1915; Kakhadze, 1942; Colo, 1961, 1964; Rieber, 1963; Elmi, 1967; Seyed-Emami, 1967; Contini, 1970; Braun and Jordan, 1976; Elmi and Caloo, 1985; Benshili, 1989; Rostovtsev, 1991; Westermann, 1992; Elmi and Rulleau, 1994; Rulleau, 1995).
3. Mediterranean (or West Tethyan) Province (shelfal or oceanic areas of North Africa, southern Spain, Italy, Austria, Hungary, Bulgaria; cf. Benecke, 1865; Vacek, 1886; Gregorio, 1886; Gemmellaro, 1886; Ramaccioni, 1934; Maubeuge, 1955; Dubar, 1960; Geczy, 1967; Fischer, 1970; Venturi, 1985; Kálin and Ureta, 1987; Mouterde and Elmi, 1991; Sadki, 1996; Callomon *et al.*, 1995; Cresta, 1996).

T. scissum has been recognized in southern Asia (cf. Sato, 1954; Sato *in* Komalarjun and Sato, 1964; Braun and Jordan, 1976) and in the West Pacific Province (Japan and far eastern of the former USSR; cf. Sato, 1954). Specimens of *T. scissum* have been found also in the following three provinces of the East-Pacific Subrealm:

1. Andean Province (western South America, Chile, Peru, Argentina; cf. Burckhardt, 1903; Westermann and Riccardi, 1972; Bogdanic *et al.*, 1985; Hillebrandt and Westermann, 1985; Hillebrandt *et al.*, 1986; Gröschke *et al.*, 1988; Riccardi *et al.*, 1990, 1992; Westermann, 1992).
2. Athabaskan Province (western North America; Frebold, 1951, 1964; Frebold *et al.*, 1969; Imlay, 1982; Poulton *et al.*, 1991; Poulton and Tipper, 1991; Poulton *et al.*, 1992).
3. Shoshonean Province (Western Interior of the United States; Imlay, 1973).

In conclusion, *T. scissum* was a pandemic species, inhabiting oceanic or shelfal environments in early Aalenian. However, some species of *Tmetoceras*, such as *T. regleyi* and

T. flexicostatum, were geographically restricted in very distant areas. *T. regleyi* has been found only in European areas of the West Tethyan Subrealm. Representatives of *T. regleyi* were distributed mainly in the Sub-Mediterranean and Northwest European provinces, inhabiting shallow environments of epicontinental platforms (cf. Dumortier, 1874; Gemmellaro, 1886; Roman and Boyer, 1923; Buckman, 1892; Schlegelmich, 1985; Rulleau, 1995). It is suggested here that a pattern of adaptive radiation may have taken place in the Western Tethys during the Opalinum-Murchisonae biochrons, giving rise to *T. regleyi* from *T. scissum*. Specialized forms of *Tmetoceras* (k-strategists such as the individuals of the species *T. regleyi*) are widespread in the epicontinental platforms around the Western Tethys during the Murchisonae and Bradfordensis biochrons. Epicontinental, specialized forms of *T. regleyi* suffered extinction in the latest Bradfordensis Biochron. Shelfal or oceanic, generalist forms of *T. scissum* disappeared in the Western Tethys or the Mediterranean Province in the latest Bradfordensis Biochron (Fernández-López *et al.*, 1997), but they survived in the East-Pacific Subrealm. In latest Bradfordensis Biochron or in earliest Concavum Biochron, *T. scissum* gave rise to *T. kirki* (Westermann), a species restricted to the East-Pacific Subrealm (Westermann, 1964, 1992; Westermann and Riccardi, 1972; Bogdanic *et al.*, 1985; Hillebrandt and Westermann, 1985; Poulton and Tipper, 1991).

From a palaeobiogeographical point of view, distinctive sedimentary basins of Iberia belonged to three adjacent bioprovinces throughout Aalenian time: Mediterranean, Sub-Mediterranean and Northwest European provinces (Figure 19). The relative orientation of the Iberian Peninsula has been accomplished on the basis of the palaeogeographical data published by Ziegler (1990) and Dercourt *et al.* (1993).

The Mediterranean Province included the Betic Basin and it occupied regions open to oceanic influence, showing characteristically high abundance of Phyllo- and Lytoceratina, where Hammatoceratidae plus Tmetoceratinae were more frequent than Graphoceratidae.

The Sub-Mediterranean Province essentially occupied the epicontinental seas bordering the northern margins of Tethys (cf. Page, 1996); in particular, eastern areas of the Iberian Basin (SE Castilian Platform, Tortosa Platform and Aragonese Platform, at least), but also some regions open to oceanic influence such as the North Lusitanian Basin. In the




Figure 19. Areas of the Iberian Peninsula occupied by three separate ammonite bioprovinces throughout Aalenian time. NW and Central areas of the Castilian platform were not inhabited by *Tmetoceras*, although some shells arrived to these shallow, epicontinental, outer platforms by nekroplanktic drift. The diversity of ammonoid faunas decrease towards the northern areas, from the Mediterranean Province to the Northwest European Province. This biogeographic gradient was latitudinally limited by environmental factors, including temperature, showing no evidences of a bathymetric control.

Sub-Mediterranean Province, endemic Phyllo- and Lytoceratina were generally absent; endemic Phyllo- and Lytoceratina were not important elements of the preserved associations; ammonoid diversity was relatively high; and Graphoceratidae were more frequent than Hammatoceratidae plus Tmetoceratinae. No endemic *Tmetoceras* are known from the NW and Central areas of the Castilian Platform.

The Northwest European Province encompassed northwestern areas of the Iberian Basin (Asturias, Basque-Cantabrian Basin and NE Cantabrias). In this province even endemic Phyllo- and Lytoceratina were generally absent; ammonoid diversity was relatively low; and Graphoceratidae representatives were predominant elements (up to 90%).

The distribution of the Sub-Mediterranean Province in the Iberian Plate (Figure 19), from areas open to oceanic influence of the Lusitanian Basin to areas occupied by shallow epicontinental seas of the Iberian Basin, allows us to exclude a bathymetric control of this bioprovince. The ammonoid provincialism through the Aalenian in these areas may have been latitudinally limited by environmental factors, including temperature rather than depth, since the diversity of ammonoid faunas decrease towards the northern areas.

5. CONCLUSIONS

Tmetoceras scissum and *T. regleyi* appear to intergrade, but they were ecological and biogeographically differentiated. In Iberia at least, populations of *T. scissum* and *T. regleyi* were not coexistent. However, by necroplanktic drift, the shells of these species may be associated in the same sites and beds.

Taphonomic data suggest a endemic character in shelfal basins or oceanic areas to the representatives of *T. scissum*. Nevertheless, exceptional immigrants and drifted shells of this species arrived in shallow environments of neighbouring platforms. In contrast, representatives of *T. regleyi* inhabited preferentially shallow environments of epicontinental platforms.

Several taphonomic criteria (type of taphonic population, abundance, frequency, stratigraphic persistence, degree of packing, and degree of taphonomic heritage) support the concept that epicontinental platforms were the sites of adaptive radiation of *T. regleyi*.

The biogeographical dispersal of *T. scissum* and *T. regleyi*, and the arrival of immigrants in the epicontinental platforms of Iberia, were favoured by the relative sea-level rise during the Opalinum and Bradfordensis biochrons. The relative sea-level fall during the Murchisonae and Concavum biochrons decreased the accessibility of *Tmetoceras* shells from shelfal basins to epicontinental platforms. The Iberian Basin was not active as an epicontinental migration route for Aalenian *Tmetoceras*.

The ammonoid provincialism throughout the Aalenian in Iberia may have been latitudinally limited by environmental factors, including temperature, since the diversity of ammonoid faunas decrease towards the northern areas.

The taphonomic interpretation of successive transients is very important in explaining the palaeobiogeographical distribution of species and the diversity of genera. The general conclusion is that taphonomic analysis of successive transients, as in this case, are relevant in interpreting the biogeographical distribution and the taxonomic diversity of ammonites.

ACKNOWLEDGMENTS

The authors wish to express their acknowledgement to Doctor E. Cariou (Univ. Poitiers) for the critical reading of the manuscript and suggestions made. The authors are

grateful to Uly Martin for the preparation of the photographs. This work was financed by the projects PB91-0383 (DGICYT), PB92-0011 (DGICYT-CSIC), PB94-0786 (DGICYT) and PB96-0838 (DGES-CSIC), and by two Luso Hispanic Integrated Actions (E9-95 and HP94-043).

REFERENCES

- Alvaro, M., Barnolas, A., Cabra, P., Comas-Rengifo, M.J., Del Olmo, P., Fernández-López, S., Goy, A., Ramírez del Pozo, J., Simo, A., and Ureta, M.S., 1989, El Jurásico de Mallorca (Islas Baleares), *Cuad. Geol. Ibérica* **13**: 67-120.
- Arkell, W.J., 1933, *The Jurassic System in Great Britain*. Clarendon Press, Oxford, pp. 1-681.
- Benecke, E.W., 1865, Über Trias und Jura in den Südalpen, *Geognostische-paläont. Beitr.* **1**: 1-204.
- Benshili, K., 1989, Lias-Dogger du Moyen Atlas plissé (Maroc), sédimentologie, biostratigraphie et évolution paléogéographique, *Doc. Géol. Fac. Sci. Lyon* **106**: 11-283.
- Bielecka, W., Birkenmajer, K., Grabowska, I., Karczewski, L., Kopik, J., Malinowska, L., Mamczar, J., Marcinkiewicz, T., Nowak, W., Orłowska-Zwolinska, T., Rogalska, M., and Styk, O., 1970, Catalogue of fossils. Jurassic: Lower and Middle Jurassic, in: *Geology of Poland*, vol. 2, part 2, Mesozoic, Wydawnictwa Geologiczne, Warsaw, pp. 23-64.
- Bogdanic, T., Hillebrandt, A. von, and Quinzio, L.A., 1985, El Aaleniano de Sierra de Varas, Cordillera de Domeyko, Antofagasta, Chile, in: *IV Congreso Geológico Chileno*, Antofagasta, pp. 58-75.
- Braun, E. von, and Jordan, R., 1976, The stratigraphy and paleontology of the Mesozoic sequence in the Mae Sot Area in Western Thailand, *Geol. Jb.* **B21**: 5-51.
- Buckman, S.S., 1887-1907, A monograph of the ammonite of the Inferior Oolite Serie, *Paleontogr. Soc.* Monogr. + Supplement, London, pp. 1-456 +1-267.
- Burckhardt, C., 1903, Beiträge zur Kenntnis der Jura- und Kreideformation der Cordillere, *Palaeontogr. Abt.* **50**: 1-144.
- Callomon, J.H., 1995, Time from fossils: S.S. Buckman and Jurassic high-resolution geochronology, in: *Milestones in Geology* (M.J. Le Bas, ed.), *Geol. Soc. London, Memoir* **16**: 127-150.
- Callomon, J.H., and Chandler, R.B., 1990, A review of the ammonite horizons of the Aalenian - Lower Bajocian Stages in the Middle Jurassic of Southern England, *Mem. Descr. Cart. Geol. Hal.* **40**: 85-112.
- Callomon, J.H., and Chandler, R.B., 1994, Some early Middle Jurassic ammonites of Tethyan affinities from the Aalenian of southern England, *Palaeopelagos*, Spec. Publ. **1**: 17-40.
- Callomon, J.H., Cresta, S., and Pavia, G., 1995, A revision of the classical Aalenian succession in the Middle Jurassic of Capo San Vigilio, northern Italy, *Geobios* M.S. **17**: 103-110.
- Canales, M.L., Goy, A., Herrero, C., and Ureta, M., 1993, Foraminíferos del Aaleniano en el sector suroccidental de la Cuenca Vasco-Cantábrica, *Treb. Mus. Geol. Barcelona* **3**: 19-40.
- Colo, G., 1961, Contribution a l'étude du Jurassique du Moyen Atlas septentrional, *Not. Mém. Serv. Géol. Maroc* **139**: 1-226.
- Colo, G., 1961-64, Contribution a l'étude du Jurassique du Moyen Atlas septentrional. Atlas des planches hors texte, *Not. Mém. Serv. Géol. Maroc* **139 bis**: 1-22.
- Contini, D., 1970, L'Aalénien et le Bajocien du Jura franc-comtois. Étude stratigraphique, *Ann. Sci. Univ. Besançon*, (3), *Geol.* **11**: 1-203.
- Cresta, S., 1996, Aalenian ammonite biostratigraphy in Northern Apennines (Italy), *GeoResearch Forum* **1-2**: 135-138.
- Dercourt, J., Ricou, L.E., and Vrielynck, B. eds., 1993, *Atlas Téthys, Palaeoenvironmental maps*, Gauthier-Villars, Paris, pp. 1-307.
- Dietl, G., 1977, The Braunjura (Brown Jurassic) in Southwest Germany, *Stuttgarter Beitr. Naturk.* **25**: 1-41.
- Dietl, G., and Etzold, A., 1977, The Aalenian at the type locality, *Stuttgarter Beitr. Naturk.* **30**: 1-13.
- Dubar, G., 1960, Supplement a l'étude des faunes aaléniennes de Krendegg et du J. Tratt (Préif, Maroc), *Ann. Soc. Géol. Nord* **80**: 50-52.
- Dumortier, E., 1874, *Études paléontologiques sur les dépôts jurassiques du Bassin du Rhône*. 4e partie, Lias supérieur, Savy, Paris, pp. 1-339.
- Elmi, S., 1967, Le Lias supérieur et le Jurassique moyen de l'Ardèche, *Doc. Lab. Géol. Fac. Sci. Lyon* **19**: 1-845.
- Elmi, S., and Caloo, P., 1985, Éléments essentiels des peuplements d'ammonites du Toarcien terminal-Aalénien en Oranie (Algérie occidentale), *Cah. Inst. Cathol. Lyon* **14**: 45-56.
- Elmi, S., and Rulleau, L., 1994, Le Jurassique du Beaujolais méridional. *Geobios* M.S. **15**: 139-155.

- Fernández-López, S., 1980, Estudio bioestratigráfico (Ammonoidea) de materiales aalenenses en Ribarroja (Valencia), *Estudios Geol.* **35** (1979): 415-423.
- Fernández-López, S., 1985, *El Bajociense en la Cordillera Ibérica*, Ph.D. Thesis. Departamento de Paleontología, Univ. Complutense Madrid, pp. 1-850.
- Fernández López, S., 1990, El significado de la autoctonía / aloctonía tafonómica, in: *Comunicaciones de la Reunión de Tafonomía y Fosilización* (S. Fernández López, coord.), Departamento Paleontología, Univ. Complutense Madrid, pp. 115-124.
- Fernández López, S., 1991a, Taphonomic concepts for a theoretical biochronology, *Rev. Esp. Paleontol.* **6**: 37-49.
- Fernández-López, S., 1995, Taphonomie et interprétation des paléoenvironnements, in: *First European Palaeontological Congress*, Lyon 1993 (M. Gayet and B. Courtinat, eds.), *Geobios M.S.* **18**: 137-154.
- Fernández-López, S., 1997, Ammonites, clinos tafonómicos y ambientes sedimentarios, *Rev. Esp. Paleontol.* **12**: 102-128.
- Fernández-López, S., and Aurell, M., 1988, El Bajociense y Bathoniense de Riela (Zaragoza). Características bioestratigráficas y sedimentológicas, *Ciencias de la Tierra*, Geología **11**: 251-264.
- Fernández-López, S., and Gómez, J.J., 1990a, Evolution tectono-sédimentaire et genèse des associations d'ammonites dans le secteur central du Bassin Ibérique (Espagne) pendant l'Aalénien, *Cah. Univ. Cathol. Lyon*, sér. sci. **4**: 39-52.
- Fernández-López, S., and Gómez, J.J., 1990b, Facies aalenenses y bajocienses, con evidencias de emersión y carstificación, en el sector central de la Cuenca Ibérica. Implicaciones paleogeográficas, *Cuad. Geol. Ibérica* **14**: 67-111.
- Fernández-López, S., and Meléndez, G., 1995, Phylloceratina ammonoids in the Iberian Basin during the Middle Jurassic. A model of biogeographical and taphonomic dispersal related to relative sea-level changes, *Palaeogeogr., Palaeoclimatol., Palaeoecol.* **120**: 291-302.
- Fernández-López, S., and Mouterde, R., 1985, Le Toarcien, l'Aalénien et le Bajocien dans le secteur de Tivenys: nouvelles données biostratigraphiques, *Strat.*, **2**: 71-82.
- Fernández-López, S., and Mouterde, R., 1994, L'Horizon à Gervillii (Bajocien inférieur) de Tendron (Cher, France). Taphonomie et populations d'ammonites. Proceedings 3rd International Meeting on Aalenian and Bajocian Stratigraphy, *Miscellanea Ser. Geol. Nazionale* **5**: 117-159.
- Fernández-López, S., and Suárez-Vega, L.C., 1980, Estudio bioestratigráfico (Ammonoidea) del Aalenense y Bajociense en Asturias, *Estudios Geol.* **35** (1979): 231-239.
- Fernández-López, S., Gómez, J.J., and Ureta, M.S., 1988a, Características de la plataforma carbonatada del Dogger en el sector meridional de la Sierra de la Demanda (Soria), *Ciencias de la Tierra*, Geología **11**: 167-195.
- Fernández-López, S., Goy, A., and Ureta, M.S., 1988b, El Toarciense superior, Aalenense y Bajociense inferior en Camino (Santander). Precisiones bioestratigráficas, *Ciencias de la Tierra*, Geología **11**: 47-62.
- Fernández-López, S., Aurell, M., García Joral, F., Gómez, J.J., Henriques, M.H., Martínez, G., Meléndez, G., and Suárez Vega, L.C., 1996, El Jurásico Medio de la Cuenca Catalana: unidades litoestratigráficas y elementos paleogeográficos, *Rev. Esp. Paleontol.* **N. extra** (1996): 122-139.
- Fernández-López, S., Henriques, M.H., Linares, A., Sandoval, J., and Ureta, M.S., 1997, Aalenian *Tmetoceras* (Ammonoidea) from Iberia: taxonomy and evolution (this volume).
- Fischer, E., 1915, Jura und Kreideversteinerungen aus Persien, *Beitr. Paläontol. Geol. Österreich-Ungarns* **27**: 207-273.
- Fischer, R., 1970, Ammoniten aus dem Aalenium der nördlichen Kalkalpen, *N. Jb. Geol. Paläontol. Mh.* **10**: 585-604.
- Flamand, G.B.M., 1911, *Recherches géologiques et géographiques sur le Haut-Pays de l'Oranie et sur le Sahara (Algérie et Territoires du Sud)*. Missions du Ministère de l'Instruction publ. et du Gouv. gen. Alg. Territoires du Sud. de l'Algérie, Serv. Géol., Lyon, pp. 1-1001.
- Frebold, H., 1951, Contributions to the palaeontology and stratigraphy of the Jurassic System in Canada, *Geol. Surv. Can. Bull.* **18**: 1-54.
- Frebold, H., 1964, Illustrations of Canadian fossils, Jurassic of Western and Arctic Canada, *Geol. Surv. Can. Pap.* **63-4**: 1-136.
- Frebold, H., Tipper, H.W., and Coates, J.A., 1969, Toarcian and Bajocian rocks and guide ammonites from southwestern British Columbia, *Geol. Surv. Can. Pap.* **67-10**: 1-55.
- García-Gómez, R., Jiménez, A.P., Linares, A., Rivas, P., and Sandoval, J., 1994, The Toarcian-Aalenian boundary in the Betic Cordillera (Southern Spain), *Geobios M.S.* **17**: 211-221.
- Geczy, B., 1967, Ammonoites jurassiques de Csernye, Montagne Bakony, Hongrie. Part II (excl. Hammatoceratidae), *Geol. Hung. (Paleontol.)* **35**: 1-413.
- Gemmellaro, G., 1886, Sul Dogger inferiore di Monte San Giuliano, *Bull. Soc. Sci. Nat. Econ. Palermo* **17**: 197-213.

- Goy, A., and Ureta, S., 1981, Bioestratigrafía del Aalenense en Préjano-Muro de Aguas (Logroño), *Cuad. Geol.* **10**: 107-119.
- Goy, A., and Ureta, S., 1987, Leioceratinae (Ammonitina) del Aalenense inferior de Fuentelsaz (Cordillera Ibérica, España), *Bull. Soc. Paleontol. Ital.* **25**: 213-236.
- Goy, A., and Ureta, S., 1988, Ammonitina del Toarciense superior en la Sierra Norte de Mallorca (España), *Bol. Real Soc. Esp. Hist. Nat. (Geol.)* **84**: 19-38.
- Goy, A., and Ureta, S., 1991, The lower boundary of the Aalenian in the Fuentelsaz section (Iberian Range, Spain), in: *Conference on Aalenian and Bajocian Stratigraphy* (N. Morton, ed.), Birkbeck College, Univ. London, pp. 33-47.
- Goy, A., Martínez, G., and Ureta, S., 1988, Bioestratigrafía del Toarciense y Aalenense en el sector comprendido entre La Almunia de Doña Godina y Ricla (Provincia de Zaragoza), *Ciencias de la Tierra*, Geología **11**: 237-250.
- Goy, A., Martínez, G., and Ureta, M.S., 1990, The Upper Toarcian and the Aalenian in the Puerto de Pozazal area (Western Basque-Cantabrian Basin, Northern Spain), in: *VI Meeting European Geological Society*, Lisboa, Abstracts: 31.
- Goy, A., Ureta, S., Arias, C., Canales, M.L., García Joral, F., Herrero, C., Martínez, G., and Perilli, N., 1994, The Fuentelsaz section (Iberian range, Spain), a possible stratotype for the base of the Aalenian stage, *Miscellanea Ser. Geol. Nazionale* **5**: 1-31.
- Gregorio, A. de, 1886, *Iconografía de la fauna dell'orizzonte Alpiniano*, (L.P. Lauriel), Palermo, pp. 1-6.
- Gröschke M., Hillebrandt A. v., Prinz P., Quinzio L.A., and Wilke H.-G., 1988, Marine Mesozoic paleogeography in Northern Chile between 21° - 26° S, *Lecture Notes Earth Sci.* **17**: 105-117.
- Henriques, M.H., 1989, O limite Lias-Dogger no Baixo Mondego (Portugal) - Ensaio de caracterização bioestratigráfica, *Memorias e Noticias* **107**: 37-42.
- Henriques, M.H., 1992, *Biostratigrafia e Paleontologia (Ammonitina) do Aaleniano em Portugal (sector setentrional da Bacia Lusitaniana)*, Ph.D. Thesis, Univ. Coimbra, pp. 1-301.
- Henriques, M.H., 1995, Les faunes d'ammonites de l'Aalénien portugais: composition et implications paléobiogéographiques, *Geobios M.S.* **18**: 229-235.
- Henriques, M.H., Linares, A., Sandoval, J., and Ureta, M.S., 1996, The Aalenian in the Iberia (Betic, Lusitanian and Iberian basins), *GeoResearch Forum* **1-2**: 139-150.
- Hernández-Molina, F.J., Sandoval, J., Aguado, R., O'Dogherty, L., Comas, M.C., and Linares, A., 1991, Olistoliths from the Middle Jurassic in the Cretaceous materials of the Fardes Formation. Biostratigraphy (Subbetic zone, Betic Cordillera), *Rev. Soc. Geol. Esp.* **4**: 79-104.
- Hillebrandt, A. von, and Westermann, G.E.G., 1985, Aalenian (Jurassic) Ammonite faunas and zones of the Southern Andes, *Zitteliana* **12**: 3-55.
- Hillebrandt, A. v., Gröschke, M., Prinz, P., and Wilke, H.G., 1986, Marines Mesozoikum in Nordchile zwischen 21° und 26° S, *Berliner geowiss. Abh.* **66**: 169-190.
- Imlay, R.W., 1955, Characteristic Jurassic mollusks from northern Alaska, *U.S. Geol. Surv. Prof. Paper* **274-D**: 69-96.
- Imlay, R.W., 1962, Late Bajocian ammonites from the Cook Inlet Region, Alaska, *U.S. Geol. Surv. Prof. Paper* **418-A**: 1-14.
- Imlay, R.W., 1964, Middle Bajocian ammonites from the Cook Inlet Region, Alaska, *U.S. Geol. Surv. Prof. Paper* **418-B**: 1-61.
- Imlay, R.W., 1973, Middle Jurassic (Bajocian) ammonites from Eastern Oregon, *U.S. Geol. Surv. Prof. Paper* **756**: 1-100.
- Imlay, R.W., 1982, Late Bajocian ammonites from Southern Alaska, *U.S. Geol. Surv. Prof. Paper* **1189**: 1-19.
- Kakhadze, J., 1942, La faune du Jurassique moyen de la Géorgie, *Bull. Inst. Geol. Géorgie* (3), **1**: 295-311.
- Kálin, O., and Ureta, S., 1987, El Lias superior y el Dogger inferior en Gorgo a Cerbara (Apenino central): aspectos bioestratigráficos y sedimentológicos, *Estudios Geol.* **43**: 489-511.
- Komalarjun, P., and Sato, T., 1964, Aalenian (Jurassic) Ammonites from Mae Sot, Northwestern Thailand, *Jap. J. Geol. Geogr.* **35**: 149-161.
- Linares, A., 1979, El Aalenense de la Zona Subbética, *Cuad. Geol.* **10**: 413-424.
- Linares, A., and Sandoval, S., 1992, El género *Vacekia* (Grammoceratinae, Ammonitina) del Aalenense de la Cordillera Bética, *Rev. Esp. Paleontol.* N. extra (1992): 91-99.
- Linares, A., and Sandoval, J., 1993, El Aalenense de la Cordillera Bética (Sur de España), análisis bioestratigráfico y caracterización paleogeográfica, *Rev. Soc. Geol. Esp.* **6**: 177-206.
- Linares, A., Ureta, M.S., and Sandoval, J., 1988, Comparison between the Aalenian ammonite association from the Betic and Iberian cordilleras: elements of correlation, in: *2nd International Symposium on Jurassic Stratigraphy* (R.B. Rocha and A.F. Soares, eds.), INIC, Lisboa, pp. 193-208.

- Maubeuge, P.L., 1955, Les Ammonites aaléniennes, bajociennes et bathoniennes du Jura Suisse septentrional. 1ère partie, *Mém. Suisses Paléontol.* **71**: 1-48.
- Morton, N., 1965, The Bearreraig Sandstone Series (Middle Jurassic) of Skye and Ransay, *Scott. J. Geol.* **1**: 189-216.
- Morton, N., and Hudson, J., 1996, Field guide to the Jurassic of the Isles of Raasay and Skye, Inner Hebrides, NW Scotland, in: *Milestones in Geology* (M.J. Le Bas, ed.), *Geol. Soc. London*, Memoir **16**: 209-280.
- Mouterde, R., and Elmi, S., 1991, Caractères différentiels des faunes d'ammonites du Toarcien des bordures de la Téthys. Signification paléogéographique, *Bull. Soc. géol. Fr.* **162**: 1185-1195.
- Mouterde, R., Rocha, R., and Ruget, Ch., 1980, Stratigraphie et faune du Lias et de la base du Dogger au Nord du Mondego (Quiaios et Brenha) (parties 2 à 4), *Comunic. Ser. Geol. Port.* **66**: 79-97.
- Mouterde, R., Ruget, Ch., and Moitinho de Almeida, F., 1964, Coupe du Lias au Sud de Condeixa, *Comunic. Ser. Geol. Port.* **48**: 61-91.
- Mouterde, R., Fernández-López, S., Goy, A., Linares, A., Rivas, P., Ruget, Ch., and Suárez-Vega, L.C., 1978, El Jurásico en la región de Obón. *Grupo Español de Mesozoico*, Jurásico Cordillera Ibérica, II: 13 p.
- Page, K.N., 1996, Mesozoic ammonoids in space and time, in: *Ammonoid Paleobiology* (N.H. Landmann, K. Tanabe and R.A. Davis, eds.), *Topics in Geobiology* **13**: 755-794.
- Patrulus, D., and Popa, E., 1971, Lower and Middle Jurassic ammonite zones in the Roumanian Carpathians, *Ann. Inst. Geol. Publ. Hung.* **54**: 131-148.
- Perrot, Ch., 1955, Études sur l'Aalénien inférieur au Nord du Tage, *Comunic. Ser. Geol. Port.* **33**: 129-151.
- Poulton, T.P., 1982, Paleogeographic and tectonic implications of the Lower and Middle Jurassic facies patterns in northern Yukon Territory and adjacent Northwest Territories, in: *Arctic Geology and Geophysics* (A.F. Embry and H.R. Balkwill, eds.), *Can. Soc. Petroleum Geol.*, Memoir **8**: 13-27.
- Poulton, T.P., and Tipper, H.W., 1991, Aalenian ammonites and strata of western Canada, *Geol. Surv. Can.* **411**: 1-71.
- Poulton, T.P., Dettmerman, R.L., Hall, R.L., Jones, D.L., Peterson, J.A., Smith, P., Taylor, D.G., Tipper, H.W., and Westernmann, G.E.G., 1992, Western Canada and United States, in: *The Jurassic of the Circum-Pacific* (G.E.G. Westernmann, ed.), Cambridge University Press, Cambridge, pp. 29-92.
- Poulton, T.P., Hall, R.L., Tipper, H.W., Cameron, B.E.B., and Carter, E.S., 1991, Current status of Middle Jurassic biostratigraphy of the Queen Charlotte Islands, British Columbia, *Geol. Surv. Can. Pap.* **90-10**: 237-252.
- Ramaccioni, G., 1939, Fauna giurassica e cretacea del Monte Cucco e dintorni (Appennino Centrale), *Palaeontol. Ital.* **39**: 143-213.
- Riccardi, A.C., Westernmann, G.E.G., and Damborenea S.E., 1990, Middle Jurassic of South America and Antarctic Peninsula, *Newsl. Stratigr.* **21**: 105-128.
- Riccardi, A.C., Gulisano, C.A., Mojica, J., Palacios, O., Schubert, C., and Thomson, M.R.A., 1992, Western South America and Antarctica, in: *The Jurassic of the Circum-Pacific* (G.E.G. Westernmann, ed.), Cambridge University Press, Cambridge, pp. 122-161.
- Rieber, H., 1963, Ammoniten und Stratigraphie des Braunjura ß der Schwäbischen Alb, *Palaeontogr. Abt.* **122**: 1-89.
- Rocha, R.B., Manuppella, G., Mouterde, R., Ruget, C., and Zbyszewski, G., 1981, Carta Geológica de Portugal 1/50.000, folha 19-C, Figueira da Foz, *Ser. Geol. Port.*, Lisboa, pp. 1-126.
- Roman, F., and Boyer, P., 1923, Sur quelques Ammonites de la Zone à "*Ludwigia murchisonae*" du Lyonnais, *Trav. Lab. Geol. Fac. Sci. Lyon* **4**: 1-47.
- Rostovtsev, K.O., 1991, On the Aalenian and Bajocian zones of the Caucasus. *Conference on Aalenian and Bajocian Stratigraphy*, in: (N. Morton, ed.), Birkbeck College, University of London, pp. 103-107.
- Rulleau, L., 1995, *Les Graphoceratidae du Toarcien supérieur et de l'Aalénien de la région Lyonnaise*, Section Géologie-Paléontologie du C.E. des Ciments Lafarge France, pp. 1-28.
- Sadki, D., 1996, Le Haut-Atlas central (Maroc): Stratigraphie et paléontologie du Lias supérieur et du Dogger Inférieur - Dynamique du bassin et des peuplements, *Docum. Lab. Geol. Lyon* **142**: 1-145.
- Sandoval, J., 1983, *Bioestratigrafía y Paleontología (Stephanoceratidae y Perisphinctidae) del Bajocense y Bathonense en las Cordilleras Béticas*, Ph.D. Thesis, Univ. Granada, pp. 1-613.
- Sato, T., 1954, Découverte de *Tmetoceras* dans le plateau de Kitakami au Nord du Japon, *Jap. J. Geol. Geogr.* **24**: 11-121.
- Schlegelmilch, R., 1985, *Die Ammoniten des süddeutschen Doggers*, Gustav Fischer Verlag, Stuttgart, pp. 1-284.
- Seyed-Emami, K., 1967, *Zur Ammoniten-Fauna und Stratigraphie der Badami-Kalke bei Kerman, Iran (Jura, oberes Untertoarcium bis mittleres Bajocium)*, Ph.D. Thesis, Ludwig Maximilians Univ. München, pp. 1-180.
- Seyfried, H., 1978, Der Subbetiche Jura von Murcia (Südost-Spanien), *Geol. Jb.* **29**: 3-201.
- Suárez Vega, L.C., 1974, Estratigrafía del Jurásico de Asturias, *Cuad. Geol. Ibérica* **3**: 1-370.

- Ureta, S., 1985, *Bioestratigrafía y paleontología (Ammonitina) del Aalenense en el sector noroccidental de la Cordillera Ibérica*, Ph.D. Thesis, Editorial Complutense, n. 158/85, Madrid, pp. 1-452.
- Ureta, S., 1988, El Aalenense en el borde nororiental de la Sierra de los Cameros (Muro de Aguas, Logroño), *Ciencias de la Tierra*, Geología **11**: 299-308.
- Ureta, S., and Goy, A., 1986, El Aalenense en el área de Talveila (Soria). Bioestratigrafía y evolución sedimentaria, *Estudios Geol.* **42**: 331-339.
- Vacek, M., 1886, Ueber die Fauna der Oolithe von Cap San Vigilio verbunden mit einer Studie ueber die obere Liassgrenze, *Abh. k.k. Geol. Reichsanst.* **12**: 57-212.
- Venturi, F., 1985, *Ammoniti liassici dell'Appennino centrale*, Perugia, pp. 1-126.
- Westermann, G.E.G., 1964, The ammonite fauna of the Kialagvik Formation at Wide Bay, Alaska Peninsula. Part I. Lower Bajocian (Aalenian), *Bull. Amer. Paleontol.* **47**: 325-462.
- Westermann, G.E.G., 1981, Ammonite biochronology and biogeography of the Circum Pacific Middle Jurassic, in: *The Ammonoidea* (M.R. House and J.R. Senior, eds.), *Systematics Association*. Spec. vol. 18, Academic Press, London, pp. 459-498.
- Westermann, G.E.G., 1992, *The Jurassic of the Circum-Pacific*. Cambridge University Press, Cambridge, pp. 1-676.
- Westermann, G.E.G., and Riccardi, A.C., 1972, Middle Jurassic ammonoid fauna and biochronology of the Argentinian-Chilean Andes. Part I: Hildocerataceae, *Palaeontogr. Abt* **140**: 1-116.
- Ziegler, P.A., 1990, *Geological Atlas of Western and Central Europe 1990*, Shell Internationale Petroleum Maatschappij, pp. 1-239.